

Apsidal Behavior Among Planetary Orbits: Testing the Planet-Planet Scattering Model

Rory Barnes¹ and Richard Greenberg¹

ABSTRACT

Planets in extrasolar systems tend to interact such that their orbits lie near a boundary between apsidal libration and circulation, a “separatrix”, with one eccentricity periodically reaching near-zero. One explanation, applied to the ν And system, assumed three original planets on circular orbits. One is ejected, leaving the other two with near-separatrix behavior. We test that model by integrating hundreds of hypothetical, unstable planetary systems that eject a planet. We find that the probability that the remaining planets exhibit near-separatrix motion is small ($< 5\%$ compared with nearly 50% of observed systems). Moreover, while observed librating systems are evenly divided between aligned and anti-aligned pericenter longitudes, the scattering model strongly favors alignment. Alternative scattering theories are proposed, which may provide a more satisfactory fit with observed systems.

Subject headings: methods: n -body simulations, stars: planetary systems, planets and satellites: formation

1. Introduction

A significant fraction ($\sim 50\%$) of adjacent pairs of planets lie near a “secular separatrix”, *i.e.* a boundary in orbital element space between apsidal libration (the difference in the longitudes of periastron $\Delta\varpi$ oscillates about a fixed value) and circulation ($\Delta\varpi$ oscillates through 360°) (Barnes & Greenberg 2006a, 2006b). One characteristic of this behavior is that one orbit periodically becomes circular. For the ν And system, an archetypal near-separatrix system, Ford *et al.* (2005) suggested a model involving an unstable system of three planets on initially coplanar, circular orbits in which a gravitational encounter ejects one planet, leaving a pair of planets still bound to the star. The event created a new “initial condition”

¹Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ 85721, rory@lpl.arizona.edu, greenber@lpl.arizona.edu

with one planet on an eccentric orbit and the other still on a nearly circular orbit, such that the subsequent secular interaction is near-separatrix.

This “planet-planet scattering” model was first posited to explain the large eccentricities of extra-solar planets (Rasio & Ford 1996). Malhotra (2002) invoked a simplified version (one planet was massless) of this model to explain what was then believed to be a high fraction of systems exhibiting apsidal libration (*e.g.* Zhou & Sun 2003).

In fact, based on improved observations and statistics, Barnes & Greenberg (2006a) found that libration is relatively rare ($\sim 15\%$ of cases). What is common, whether a system librates or circulates, is to be near the boundary between those states. Ford *et al.* (2005) described only one specific hypothetical case that, when integrated numerically, did result in two planets near a secular separatrix. However, that case is anecdotal. Here we consider whether simulations like those in the Ford *et al.* model of planet-planet scattering can statistically reproduce the observed large fraction of systems that lie near the secular separatrix, as well as the observed distribution among circulating, aligned librating and anti-aligned librating systems (which Ford *et al.* does not address). Here we systematically survey hundreds of initial conditions, similar to the case described in Ford *et al.*, in order to consider whether planet-planet scattering can explain the characteristics of the observed systems.

2. Methodology

We consider a hypothetical system of a $1.3 M_{\odot}$ star and three planets, called 1, 2, and 3, with respective semi-major axes of 0.83, 3.555 and 4.4 AU, and masses of 1.94, 3.94 and 1.32 Jupiter masses (properties similar to the observed *v* And system and the hypothetical configuration considered in Ford *et al.*). All these orbits are circular and coplanar. We choose an initial condition with planet 1 45° ahead of planet 3 in longitude, L . We then consider 360 similar cases, but with the initial longitude of planet 2, L_2 , distributed evenly around 360° in 1° intervals. With these masses and orbits, the outer two planets fail a known stability condition (Gladman 1993; Barnes & Greenberg 2006c), which is independent of L . We also considered a sampling of cases with different semi-major axes and masses.

We use the symplectic, N -body integrator MERCURY (Chambers 1999) to integrate each case for 10^5 years. We require each simulation to conserve energy to within 1 part in 10^4 , which has been shown to be sufficiently accurate for symplectic integration methods (Barnes & Quinn 2004). Our smallest timestep was 10^{-3} days. For this level of energy conservation, angular momentum conservation was always at least 1 order of magnitude better. For configurations that eject the outer planet, we then integrate the remaining

planets for 10^5 years in order to characterize the secular interaction of those remaining planets, *i.e.* to calculate the orbit’s proximity to the separatrix and to determine the type of apsidal interaction. In a few cases we integrated for 5×10^5 years because the resultant secular period was longer than 10^5 years.

3. Results

After 100,000 years, of the sample of 360 cases with varying L_2 , 169 ejected the outer planet leaving two planets engaged in ongoing secular interactions that could be characterized in meaningful ways. An additional 95 cases also ejected the outer planet, but left one of the remaining planets with $a > 6$ AU, too far to be observed by current search methods. Of the remaining cases, 37 ejected no planets, but left all of them on highly eccentric orbits that interacted in complex and chaotic ways. Another 5 ejected only the middle planet, again leaving planets in highly eccentric, unstable, chaotically interacting orbits. In 49 cases two planets were ejected. Finally, 5 cases were rejected on the technical grounds that energy was not conserved to a level that could guarantee our desired precision. This number of cases is too small to affect the resulting statistics.

In order to have an outcome with near-separatrix motion, as envisioned for example in the scenario proposed by Ford *et al.*, a certain sequence of events appears to be required. Fig. 1a shows the evolution of one of our cases that does yield such an outcome. First, within only a few years, an interaction between planets 2 and 3 yields $e_3 > 0.7$ and a substantial increase in e_2 . Planet 3 spends most of its time far from the other planets, so the inner two planets undergo secular interactions independent of the third. Because e_2 becomes non-zero while the inner orbit remains circular, their secular behavior is characterized by periodic returns of e_1 to zero and typical near-separatrix behavior. Eventually, the outer planet might have a close encounter with one or both of the inner planets, which would wreak havoc with the regular secular behavior. However, before that can happen, within a few thousand years, planet 3 receives a small kick that ejects it from the system. The kick required to eject planet 3 is small enough that it does not significantly affect the secular interaction of 1 and 2, and their near-separatrix behavior is preserved.

Thus the requirements for near-secular behavior seem to be (1) a quick large increase in e_3 , with a modest increase in e_2 while e_1 remains zero, followed by (2) an encounter that ejects planet 3 without disturbing the other planets too much.




Fig. 1.— Evolution of orbital eccentricities in four characteristic cases. (a) This case, with the initial longitude of planet 2 relative to planet 3 of 123° , shows a typical series of events leading to near-separatrix motion. (b) In this case, the initial encounter starts near-separatrix behavior of planets 1 and 2, but planet 3 remains in the system and disrupts the regular secular behavior $L_2 = 274^\circ$. (c) Here, the initial encounter fails to increase e_3 enough to uncouple it from the other planets ($L_2 = 90^\circ$). (d) An adequate early jump in e_3 , is soon followed by disruptive interactions that eject planet 3. However the remaining planets' secular interaction ultimately results in a destabilizing encounter ($L_2 = 314^\circ$).

Fig. 1b shows a case that satisfies the first condition, but not the second. Planet 3 quickly enters a highly eccentric orbit, and the other planets begin to behave like a near-separatrix case, but planet 3 isn’t immediately ejected. By 10,000 years it begins to disrupt the regular two-planet secular behavior of the other planets, allowing them to evolve onto orbits where e_1 never returns to zero, *i.e.* no longer near a separatrix.

Fig. 1c shows what can happen if the first requirement is not met. Here the initial increase in e_3 is too little to keep it out of the way of the other planets. This example is one of the 37 cases that left all three planets interacting in ways that preclude regular secular behavior.

Fig. 1d shows a case where planet 3 does satisfy the first requirement, and it is also ejected from the system fairly quickly (in $\sim 10,000$ yr as in Fig. 1a), but before being ejected it has encountered and grossly de-circularized the other two orbits. After planet 3 is ejected, planets 1 and 2 begin a secular interaction, but this interaction leads to large values of e_1 and an encounter that ejects planet 1.

Next we characterize the outcomes of our cases for comparison with Ford *et al.*’s suggestion that this process can explain behavior like that of the near-separatrix ν And system, and also for comparison with the more general statistics of observed systems. A way to quantify how close a given system is to a separatrix was introduced by Barnes and Greenberg (2006a). Loosely described, a parameter ϵ represents the ratio between the minimum e value and the amplitude of oscillation of e (see Barnes and Greenberg [2006a] for a precise definition). Small ϵ means the system is near-separatrix.

Figure 2 shows the distribution of ϵ values as a fraction of the 360 initial cases. Only the 169 cases that produced regular secular behavior contribute to these statistics; for the remainder of the cases ϵ would be meaningless and near-separatrix motion is out of the question. The distribution shows a very slight rise for small values of ϵ : About 4% have ϵ smaller than 0.01 and 12% have values less than 0.03. We also show the distribution for the subset of cases in which the outer planet was ejected in only 20,000 years, with a similar distribution.

All simulations that resulted in $\epsilon < 0.01$ had final a_2 (semi-major axis of planet 2) in the range 2.84 ± 0.04 AU. Overall, 30% of simulations ended with a_2 in this range.

In addition to varying the initial longitude, we performed 8 integrations in which a_2 was varied by 0.01 AU, and 8 which varied the mass of planet 2 by 0.01 Jupiter masses. Planet-planet scattering in these cases also resulted in 1 configuration near the secular separatrix, consistent with Fig. 2.




Fig. 2.— Distribution of ϵ , a parameter which describes how close a system is to a separatrix. Squares represent the distribution for observed systems. The solid line shows statistics for our calculated outcomes of the planet-planet scattering model proposed by Ford *et al.* (2005). The real systems tend to be much closer to the separatrix (small ϵ) than can be reproduced by the Ford *et al.* hypothesis. The dashed line shows the subset of the modeled cases where ejection of planet 3 occurred within 20,000 yr. The bin size for these histograms is 0.01.

The distribution of ϵ values among actual observed planetary systems is much more concentrated near the separatrix, with very small ϵ values, as reported by Barnes and Greenberg (2006a). In Fig. 2 we include the statistics for the observed systems for comparison with the results generated by the planet-planet scattering model. For the observed values of ϵ , we use the results from Barnes and Greenberg (2006a), plus values of ϵ calculated for the two recently discovered systems HIP 14810 (Wright *et al.* 2007) and HD 160691 (Pepe *et al.* 2007), see the Appendix. Note that e is poorly known for 47 UMa and GJ 876 (Butler *et al.* 2006), but the statistics are not affected by these two systems. Similarly, the result would be unchanged if we excluded resonant and/or tidally-evolved systems (Barnes & Greenberg 2006a). As shown in Fig 2, in the observed distribution, 45% of the systems have ϵ smaller than 0.01. (In fact, the results for HIP 14810 and HD 160691 in the Appendix show the same distribution; two of the four pairs evolve with $\epsilon < 0.01$.)

Based on our numerical experiments, the planet-planet scattering model as described by Ford *et al.* (2005) does not seem to reproduce the observed strong tendency for systems to lie near a separatrix. While selected orbital configurations can lead to a small- ϵ system like v And, in general this model does not appear to reproduce that large fraction of systems that exhibit behavior very near a secular separatrix.

Of course, our experiments sample only a small portion of the possible multi-dimensional parameter space, but it gives a more reliable estimate of the probability of outcomes than the single case shown in Ford *et al.* Moreover, our tests show one specific sequence of events (*e.g.* Fig. 1a) leads to near-separatrix motion in the manner envisioned by Ford *et al.* It would be surprising if such a sequence were common, given the results presented here. Other sequences may be possible, but they are not present in our simulations. Further tests would be useful, but it seems unlikely that the planet-planet scattering model of Ford *et al.* can explain the preponderance of near-separatrix systems.

Another disagreement between the statistical outcomes of the planet-planet scattering model and actual systems comes from consideration of the modes of libration. In secular interactions, orbits can librate about an alignment of their major axes with their pericenters at the same longitude or an anti-alignment where the pericenters are 180° apart. Among observed systems (again based on calculations from Barnes and Greenberg [2006a] and the Appendix herein), there is nearly equal division between systems librating about aligned pericenters (2 out of 31) and those librating about anti-aligned pericenters (also 2 out of 31), with the remaining 87% circulating rather than librating. Among the systems generated by the planet-planet scattering model, librating cases overwhelmingly favor aligned pericenters (by a factor > 10) relative to anti-alignment. Specifically, 24% are aligned and less than 2% are anti-aligned. The planet-planet scattering model does not produce systems consistent

with the population of observed planets.

4. Conclusions

Our numerical experiments demonstrate that the planet-planet scattering process, as described by Ford *et al.* (2005), does not explain the prevalence of near-separatrix apsidal behavior, even though the model had been proposed to explain this type of motion in the case of ν And. That model can yield near-separatrix behavior, but the probability is too low for it to be a significant factor. The model does produce systems consistent with real systems in one respect: Most of them undergo apsidal circulation rather than libration. However, it produces about twice as many librating cases as are observed in reality, and it yields far too small a portion of those in anti-aligned libration compared with observed systems.

Another problem with Ford *et al.*'s planet-planet scattering is that the initial set up involves planets on circular orbits that are too close to be stable. It is difficult to envision a formation process that could yield such a system. For example, if a hypothetical eccentricity-damping medium were invoked to explain the circular orbits, the medium would need to disappear in less than a synodic period to provide the initial condition in the planet-planet scattering scenario. Even if an explanation can be found for such an initial set-up, our experiments suggest that the statistics of observed behavior would not follow.

We propose a modification to the scattering model that may explain the propensity for producing near-separatrix orbits based on conditions consistent with other lines of evidence about the origins of planetary systems. From our simulations of scattering in this current study, we find that the key features of those scattering events that lead to near-separatrix behavior are an abrupt, modest increase in the eccentricity of one planet, while another planet remains on a circular orbit, followed by a rapid removal of the cause of the perturbation (a third planet) from the system. The problems with starting the perturbing planet on a circular orbit are (1) that it is hard to understand why an unstable orbit would be circular (as mentioned above) and (2) after the initial encounter, this planet is rarely ejected from the system soon enough to keep from further modifying the interaction of the other planets.

Suppose instead that the perturbing planet started not on an unstable circular orbit, but rather on a long-period, high-eccentricity orbit. Such high-eccentricity bodies scattered about the solar system have been invoked to explain basic properties of the system, including the origin of the Moon by an impact into the Earth, pumping the relative velocities among asteroids to suppress planet growth in that zone, and generating the late heavy bombardment (Gomes *et al.* 2005; Tsiganis *et al.* 2005). Indeed the temporary passage of large bodies

scattered from the outer solar system, at a time when the inner planets had achieved fairly stable, near-circular orbits is a standard component of current models of the formation of our solar system (*e.g.* Gomes *et al.* 2005; Strom *et al.* 2006). We call this model the Rogue Planet Model.

As such a rogue planet or protoplanet passed through the inner part of a planetary system, eventually it would pass close enough to one of the regular (circularly orbiting) planets to impose a velocity change and introduce some orbital eccentricity. At the same time, because the impulsive perturber was on an extended orbit from the outer part of the system, there would be a substantial probability that it would be ejected from the system by the same encounter, preventing any further impulses on inner-system planets. After the encounter, any other regular planet will still be on a near-circular orbit, so that the subsequent secular behavior will be near-separatrix.

In order to test our proposed modification of the planet scattering model, the hypothesis should be explored with numerical experiments, analogous to those presented here and in Gomes *et al.* (2005). In our hypothesis, the prevalence of near-separatrix behavior in extra-solar planetary systems, as well as many of the dynamical properties of our Solar System, can be described by late-stage scattering of protoplanets. If these bodies are scattered inward, toward planets on circular, stable orbits, the protoplanets are ejected, leaving planets on near-separatrix orbits. If scattered out, they may become the cores of ice giants, or become part of a scattered disk component of a Kuiper Belt. If this hypothesis is correct, then the origin of the Solar System’s small eccentricities, extra-solar planets’ large eccentricities, and all planetary systems’ tendency to lie near a secular separatrix is explained by a single model.

This work was funded by NASA’s Planetary Geology and Geophysics program grant NNG05GH65G. We thank Fred Rasio, Eric Ford and an anonymous referee for useful suggestions that clarified this manuscript.

REFERENCES

- Barnes, R. & Greenberg, R. 2006a, ApJ, 652, L53
 ———. 2006b, ApJ, 638, 478
 ———. 2006c, ApJ, 647, L163
 Barnes, R. & Quinn, T.R. 2004, ApJ, 611, 494
 Butler, R.P. *et al.* 2006, ApJ, 646, 505
 Chambers, J., 1999, MNRAS, 304, 793
 Ford, E.B., Lystad, V., & Rasio, F.A. 2005, Nature, 434, 873
 Gladman, B. 1993, Icarus, 106, 247

- Gomes, R. *et al.* 2005, *Nature*, 435, 466
 Malhotra, R. 2002, *ApJ*, 575, L33
 Pepe, F. *et al.* 2007, *A&A*, 462, 769
 Rasio, F.A., & Ford, E.B. 1996, *Science*, 274, 954
 Strom, R.G. *et al.* 2006, *Science*, 309, 1847
 Tsiganis *et al.* 2005, *Nature*, 435, 459
 Wright, J.T. *et al.* 2007, *ApJ* in press
 Zhou, J.-L., & Sun, Y.-S. 2003, *ApJ*, 598, 1290

Appendix

Since the publication of Barnes & Greenberg (2006a), two planetary systems have been announced or revised. HIP 14810 (Wright *et al.* 2007) has two planetary mass companions, one of which has been tidally circularized. HD 160691, also called μ Ara, now has four planets (Pepe *et al.* 2007), of which the innermost is also tidally evolved. HD 160691 is unstable over long timescales ($\sim 10^8$ years), and its properties are therefore especially suspect. We have performed a dynamical analysis of the best-fit orbits to these two systems in the same manner as Barnes & Greenberg (2006a) in order to calculate the apsidal behavior and ϵ . These properties are listed in Table 1, where C stands for circulation and L_{180} for anti-aligned libration. The designation C/C, as in Barnes & Greenberg (2006a), means the system lies near a “circulation mode separatrix”.

Table 1: Apsidal Properties of HIP 14810 and HD 160691

System	Pair	ϵ	Apsidal Behavior
HIP 14810	b-c	0.05	L_{180}
HD 160691	c-d	0.002	C/C
	d-b	0.003	C/C
	b-e	0.13	C